


FORM PTO-1449 U.S. Department of Commerce Patent and Trademark Office				Attorney Docket Number 5470-255			Serial No. 09/541,462
LIST OF DOCUMENTS CITED BY APPLICANT (Use several sheets if necessary)				RECEIVED AUG 21 2000			
				Applicants: Xue Xiong, Tomohiko Ohta TECH CENTER 1600 Group 16430			
				Filing Date March 31, 2000			
U. S. PATENT DOCUMENTS 1652							
Examiner Initial		Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate
<i>JR</i>	1.	5,871,973	2/16/99	Hillman et al.	435	69.1	
<i>JR</i>	2.	5,922,318	7/13/99	Bandman et al.	424	94.1	
	3.	5,968,747	10/19/99	Hillman et al.	435	6	
	4.	5,968,761	10/19/99	Rolfe et al.	435	15	
	5.	5,968,797	10/19/99	Ni et al.	435	193	
<i>JR</i>	6.	6,068,982	5/30/00	Rolfe et al.	435	7.21	
<i>JR</i>	7.	6,068,994	5/30/00	Barr	435	69.7	
FOREIGN PATENT DOCUMENTS							
		Document Number	Date	Country	Class	Subclass	Translation Yes No
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)							
<i>JR</i>	8.	O. Cohen-Fix et al.; <i>Anaphase initiation in Saccharomyces cerevisiae is controlled by the APC-dependent degradation of the anaphase inhibitor Pds1p</i> ; <i>Genes & Dev.</i> 10 :3081-3093 (1996)					
<i>JR</i>	9.	A. Hershko; <i>Roles of ubiquitin-mediated proteolysis in cell cycle control</i> ; <i>Curr. Opin. Cell. Biol.</i> 9 :788-799 (1997).					
<i>JR</i>	10.	M. Brandeis et al.; <i>The proteolysis of mitotic cyclins in mammalian cells persists from the end of mitosis until the onset of S phase</i> ; <i>EMBO J.</i> 15 :5280-5289 (1996).					
<i>JR</i>	11.	N. Mathias et al.; <i>Cdc53p Acts in Concert with Cdc4p and Cdc34p To Control the G₁-to-S-Phase Transition and Identifies a Conserved Family of Proteins</i> ; <i>Mol. Cell. Biol.</i> 16 :6634-6643 (1996).					
<i>JR</i>	12.	K. M. Lonergan et al.; <i>Regulation of Hypoxia-Inducible mRNAs by the von Hippel-Lindau Tumor Suppressor Protein Requires Binding to Complexes Containing Elongins B/C and Cul2</i> ; <i>Mol. Cell. Biol.</i> 18 :732-741 (1998).					
<i>JR</i>	13.	L-C Chen et al.; <i>The Human Homologue for the <i>Caenorhabditis elegans</i> cul-4 Gene is Amplified and Overexpressed in Primary Breast Cancers</i> ; <i>Cancer Res.</i> 58 :3677-3683 (1998).					

EXAMINER
EXAMINER*JR*

DATE CONSIDERED

4/25/03

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 U.S. Department of Commerce Patent and Trademark Office		Attorney Docket Number 5470-255	Serial No. 09/541,462																																																						
LIST OF DOCUMENTS CITED BY APPLICANT																																																									
(Use several sheets if necessary)																																																									
																																																									
Applicants: Xue Xiong, Tomohiko Ohta																																																									
Filing Date March 31, 2000 Group 1643-1652																																																									
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 5%; text-align: center;"><i>DR</i></td> <td style="width: 5%; text-align: center;">14.</td> <td>J. Michel et al.; <i>Human CUL-1, but not Other Cullin Family Members, Selectively Interacts with SKP1 to Form a Complex with SKP2 and Cyclin A</i>; <i>Cell Growth Differ.</i> 9:435-449 (1998).</td> </tr> <tr> <td style="text-align: center;">↑</td> <td style="text-align: center;">15.</td> <td>M. Scheffner et al.; <i>Protein ubiquitination involving an E-1-E2-E3 enzyme ubiquitin thioester cascade</i>; <i>Nature</i> 373:81-83 (1995).</td> </tr> <tr> <td></td> <td style="text-align: center;">16.</td> <td>C. Michaelis et al.; <i>Cohesins: Chromosomal Proteins that Prevent Premature Separation of Sister Chromatids</i>; <i>Cell</i> 91:35-45 (1997).</td> </tr> <tr> <td></td> <td style="text-align: center;">17.</td> <td>R.M.R. Feldman et al.; <i>A Complex of Cdc4p, Skp1p, and Cdc53p/Cullin Catalyzes Ubiquitination of the Phosphorylated CDK Inhibitor Sic1p</i>; <i>Cell</i> 91:221-230 (1997).</td> </tr> <tr> <td></td> <td style="text-align: center;">18.</td> <td>C. Bai et al.; <i>SKP1 Connects Cell Cycle Regulators to the Ubiquitin Proteolysis Machinery through a Novel Motif, the F-Box</i>; <i>Cell</i> 86:263-274 (1996).</td> </tr> <tr> <td></td> <td style="text-align: center;">19.</td> <td>A. R. Willems et al.; <i>Cdc53 Targets Phosphorylated G1 Cyclins for Degradation by the Ubiquitin Proteolytic Pathway</i>; <i>Cell</i> 86:453-463 (1996).</td> </tr> <tr> <td></td> <td style="text-align: center;">20.</td> <td>D. Skowyra et al.; <i>F-Box Proteins Are Receptors that Recruit Phosphorylated Substrates to the SCF Ubiquitin-Ligase Complex</i>; <i>Cell</i> 91:209-219 (1997).</td> </tr> <tr> <td></td> <td style="text-align: center;">21.</td> <td>E. T. Kipreos et al.; <i>cul-1 Is Required for Cell Cycle Exit in C. elegans and Identifies a Novel Gene Family</i>; <i>Cell</i> 85:829-839 (1996).</td> </tr> <tr> <td></td> <td style="text-align: center;">22.</td> <td>A. Amon et al.; <i>Closing the Cell Cycle Circle in Yeast: G2 Cyclin Proteolysis Initiated at Mitosis Persists until the Activation of G1 Cyclins in the Next Cycle</i>; <i>Cell</i> 77:1037-1050 (1994).</td> </tr> <tr> <td></td> <td style="text-align: center;">23.</td> <td>M. Schwab et al.; <i>Yeast Hct1 Is a Regulator of Clb2 Cyclin Proteolysis</i>; <i>Cell</i> 90:683-693 (1997).</td> </tr> <tr> <td></td> <td style="text-align: center;">24.</td> <td>S. J. Sigrist et al.; <i>Drosophila fizzy-related Down-Regulates Mitotic Cyclins and is Required for Cell Proliferation Arrest and Entry into Endocycles</i>; <i>Cell</i> 90:671-681 (1997).</td> </tr> <tr> <td></td> <td style="text-align: center;">25.</td> <td>M. Scheffner et al.; <i>The HPV-16 E6 and E6-AP Complex Functions as a Ubiquitin-Protein Ligase in the Ubiquitination of p53</i>; <i>Cell</i> 75:495-505 (1993).</td> </tr> <tr> <td></td> <td style="text-align: center;">26.</td> <td>H. Funabiki et al.; <i>Cut2 proteolysis required for sister-chromatid separation in fission yeast</i>; <i>Nature</i> 381:438-441 (1996).</td> </tr> <tr> <td></td> <td style="text-align: center;">27.</td> <td>M. Glotzer et al.; <i>Cyclin is degraded by the ubiquitin pathway</i>; <i>Nature</i> 349:132-138 (1991).</td> </tr> <tr> <td></td> <td style="text-align: center;">28.</td> <td>R. W. King et al.; <i>How Proteolysis Drives the Cell Cycle</i>; <i>Science</i> 274:1652-1659 (1996).</td> </tr> <tr> <td></td> <td style="text-align: center;">29.</td> <td>J-M. Peters et al.; <i>Identification of BIME as a Subunit of the Anaphase-Promoting Complex</i>; <i>Science</i> 274:1199-1201 (1996).</td> </tr> <tr> <td style="text-align: center;">↓</td> <td style="text-align: center;">30.</td> <td>Z-K Yu et al.; <i>Human CUL-1 associates with the SKP1/SKP2 complex and regulates p21^{CIP1/WAF1} and cyclin D proteins</i>; <i>Proc. Natl. Acad. Sci. USA</i> 95:11324-11329 (1998).</td> </tr> <tr> <td style="text-align: center;"><i>DR</i></td> <td style="text-align: center;">31.</td> <td>S. A. Lyapina et al.; <i>Human CUL1 forms an evolutionarily conserved ubiquitin ligase complex (SCF) with SKP1 and an F-box protein</i>, <i>Proc. Natl. Acad. Sci. USA</i> 95:7451-7456 (1998).</td> </tr> </table>				<i>DR</i>	14.	J. Michel et al.; <i>Human CUL-1, but not Other Cullin Family Members, Selectively Interacts with SKP1 to Form a Complex with SKP2 and Cyclin A</i> ; <i>Cell Growth Differ.</i> 9 :435-449 (1998).	↑	15.	M. Scheffner et al.; <i>Protein ubiquitination involving an E-1-E2-E3 enzyme ubiquitin thioester cascade</i> ; <i>Nature</i> 373 :81-83 (1995).		16.	C. Michaelis et al.; <i>Cohesins: Chromosomal Proteins that Prevent Premature Separation of Sister Chromatids</i> ; <i>Cell</i> 91 :35-45 (1997).		17.	R.M.R. Feldman et al.; <i>A Complex of Cdc4p, Skp1p, and Cdc53p/Cullin Catalyzes Ubiquitination of the Phosphorylated CDK Inhibitor Sic1p</i> ; <i>Cell</i> 91 :221-230 (1997).		18.	C. Bai et al.; <i>SKP1 Connects Cell Cycle Regulators to the Ubiquitin Proteolysis Machinery through a Novel Motif, the F-Box</i> ; <i>Cell</i> 86 :263-274 (1996).		19.	A. R. Willems et al.; <i>Cdc53 Targets Phosphorylated G1 Cyclins for Degradation by the Ubiquitin Proteolytic Pathway</i> ; <i>Cell</i> 86 :453-463 (1996).		20.	D. Skowyra et al.; <i>F-Box Proteins Are Receptors that Recruit Phosphorylated Substrates to the SCF Ubiquitin-Ligase Complex</i> ; <i>Cell</i> 91 :209-219 (1997).		21.	E. T. Kipreos et al.; <i>cul-1 Is Required for Cell Cycle Exit in C. elegans and Identifies a Novel Gene Family</i> ; <i>Cell</i> 85 :829-839 (1996).		22.	A. Amon et al.; <i>Closing the Cell Cycle Circle in Yeast: G2 Cyclin Proteolysis Initiated at Mitosis Persists until the Activation of G1 Cyclins in the Next Cycle</i> ; <i>Cell</i> 77 :1037-1050 (1994).		23.	M. Schwab et al.; <i>Yeast Hct1 Is a Regulator of Clb2 Cyclin Proteolysis</i> ; <i>Cell</i> 90 :683-693 (1997).		24.	S. J. Sigrist et al.; <i>Drosophila fizzy-related Down-Regulates Mitotic Cyclins and is Required for Cell Proliferation Arrest and Entry into Endocycles</i> ; <i>Cell</i> 90 :671-681 (1997).		25.	M. Scheffner et al.; <i>The HPV-16 E6 and E6-AP Complex Functions as a Ubiquitin-Protein Ligase in the Ubiquitination of p53</i> ; <i>Cell</i> 75 :495-505 (1993).		26.	H. Funabiki et al.; <i>Cut2 proteolysis required for sister-chromatid separation in fission yeast</i> ; <i>Nature</i> 381 :438-441 (1996).		27.	M. Glotzer et al.; <i>Cyclin is degraded by the ubiquitin pathway</i> ; <i>Nature</i> 349 :132-138 (1991).		28.	R. W. King et al.; <i>How Proteolysis Drives the Cell Cycle</i> ; <i>Science</i> 274 :1652-1659 (1996).		29.	J-M. Peters et al.; <i>Identification of BIME as a Subunit of the Anaphase-Promoting Complex</i> ; <i>Science</i> 274 :1199-1201 (1996).	↓	30.	Z-K Yu et al.; <i>Human CUL-1 associates with the SKP1/SKP2 complex and regulates p21^{CIP1/WAF1} and cyclin D proteins</i> ; <i>Proc. Natl. Acad. Sci. USA</i> 95 :11324-11329 (1998).	<i>DR</i>	31.	S. A. Lyapina et al.; <i>Human CUL1 forms an evolutionarily conserved ubiquitin ligase complex (SCF) with SKP1 and an F-box protein</i> , <i>Proc. Natl. Acad. Sci. USA</i> 95 :7451-7456 (1998).
<i>DR</i>	14.	J. Michel et al.; <i>Human CUL-1, but not Other Cullin Family Members, Selectively Interacts with SKP1 to Form a Complex with SKP2 and Cyclin A</i> ; <i>Cell Growth Differ.</i> 9 :435-449 (1998).																																																							
↑	15.	M. Scheffner et al.; <i>Protein ubiquitination involving an E-1-E2-E3 enzyme ubiquitin thioester cascade</i> ; <i>Nature</i> 373 :81-83 (1995).																																																							
	16.	C. Michaelis et al.; <i>Cohesins: Chromosomal Proteins that Prevent Premature Separation of Sister Chromatids</i> ; <i>Cell</i> 91 :35-45 (1997).																																																							
	17.	R.M.R. Feldman et al.; <i>A Complex of Cdc4p, Skp1p, and Cdc53p/Cullin Catalyzes Ubiquitination of the Phosphorylated CDK Inhibitor Sic1p</i> ; <i>Cell</i> 91 :221-230 (1997).																																																							
	18.	C. Bai et al.; <i>SKP1 Connects Cell Cycle Regulators to the Ubiquitin Proteolysis Machinery through a Novel Motif, the F-Box</i> ; <i>Cell</i> 86 :263-274 (1996).																																																							
	19.	A. R. Willems et al.; <i>Cdc53 Targets Phosphorylated G1 Cyclins for Degradation by the Ubiquitin Proteolytic Pathway</i> ; <i>Cell</i> 86 :453-463 (1996).																																																							
	20.	D. Skowyra et al.; <i>F-Box Proteins Are Receptors that Recruit Phosphorylated Substrates to the SCF Ubiquitin-Ligase Complex</i> ; <i>Cell</i> 91 :209-219 (1997).																																																							
	21.	E. T. Kipreos et al.; <i>cul-1 Is Required for Cell Cycle Exit in C. elegans and Identifies a Novel Gene Family</i> ; <i>Cell</i> 85 :829-839 (1996).																																																							
	22.	A. Amon et al.; <i>Closing the Cell Cycle Circle in Yeast: G2 Cyclin Proteolysis Initiated at Mitosis Persists until the Activation of G1 Cyclins in the Next Cycle</i> ; <i>Cell</i> 77 :1037-1050 (1994).																																																							
	23.	M. Schwab et al.; <i>Yeast Hct1 Is a Regulator of Clb2 Cyclin Proteolysis</i> ; <i>Cell</i> 90 :683-693 (1997).																																																							
	24.	S. J. Sigrist et al.; <i>Drosophila fizzy-related Down-Regulates Mitotic Cyclins and is Required for Cell Proliferation Arrest and Entry into Endocycles</i> ; <i>Cell</i> 90 :671-681 (1997).																																																							
	25.	M. Scheffner et al.; <i>The HPV-16 E6 and E6-AP Complex Functions as a Ubiquitin-Protein Ligase in the Ubiquitination of p53</i> ; <i>Cell</i> 75 :495-505 (1993).																																																							
	26.	H. Funabiki et al.; <i>Cut2 proteolysis required for sister-chromatid separation in fission yeast</i> ; <i>Nature</i> 381 :438-441 (1996).																																																							
	27.	M. Glotzer et al.; <i>Cyclin is degraded by the ubiquitin pathway</i> ; <i>Nature</i> 349 :132-138 (1991).																																																							
	28.	R. W. King et al.; <i>How Proteolysis Drives the Cell Cycle</i> ; <i>Science</i> 274 :1652-1659 (1996).																																																							
	29.	J-M. Peters et al.; <i>Identification of BIME as a Subunit of the Anaphase-Promoting Complex</i> ; <i>Science</i> 274 :1199-1201 (1996).																																																							
↓	30.	Z-K Yu et al.; <i>Human CUL-1 associates with the SKP1/SKP2 complex and regulates p21^{CIP1/WAF1} and cyclin D proteins</i> ; <i>Proc. Natl. Acad. Sci. USA</i> 95 :11324-11329 (1998).																																																							
<i>DR</i>	31.	S. A. Lyapina et al.; <i>Human CUL1 forms an evolutionarily conserved ubiquitin ligase complex (SCF) with SKP1 and an F-box protein</i> , <i>Proc. Natl. Acad. Sci. USA</i> 95 :7451-7456 (1998).																																																							

EXAMINER
EXAMINER*BL*


DATE CONSIDERED

4/25/03

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 U.S. Department of Commerce Patent and Trademark Office		Attorney Docket Number 5470-255	Serial No. 09/541,462
LIST OF DOCUMENTS CITED BY APPLICANT			
(Use several sheets if necessary)			
 RECEIVED AUG 21 2000 TECH CENTER 1600 Group 1632			
DR	32.	S. Lahav-Baratz et al.; <i>Reversible phosphorylation controls the activity of cyclin-ubiquitin ligase</i> ; Proc. Natl Acad. Sci. USA 92 :9303-9307 (1995).	
↑	33.	Y-L Juang et al.; <i>APC-Mediated Proteolysis of Ase1 and the Morphogenesis of the Mitotic Spindle</i> ; Science 275 :1311-1314 (1997).	
	34.	R. Visintin et al.; <i>CDC20 and CDH1: A Family of Substrate-Specific Activators of APC-Dependent Proteolysis</i> ; Science 278 :460-463 (1997).	
	35.	R. Verma et al.; <i>Phosphorylation of Sic1p by G₁ Cdk Required for its Degradation and Entry into S Phase</i> ; Science 278 :455-460 (1997).	
	36.	H. Yu et al.; <i>Identification of a Cullin Homology Region in a Subunit of the Anaphase-Promoting Complex</i> ; Science 279 :1219-1222 (1998).	
	37.	W. Zachariae et al.; <i>Mass Spectrometric Analysis of the Anaphase-Promoting Complex from Yeast: Identification of a Subunit Related to Cullins</i> ; Science 279 :1216-1219 (1998).	
	38.	J. M. Huibregtse et al.; <i>A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase</i> ; Proc. Natl. Acad. Sci. USA 92 :2563-2567 (1995).	
	39.	A. Varshavsky; <i>The N-end rule: Functions, mysteries, uses</i> ; Proc. Natl. Acad. Sci. USA 93 :12142-12149 (1996).	
	40.	J. Lisztwan et al.; <i>Association of human CUL-1 and ubiquitin-conjugating enzyme CDC34 with the F-box protein p45^{SKP2}: evidence for evolutionary conservation in the subunit composition of the CDC34-SCF pathway</i> ; The EMBO J. 17 , No. 2:368-383 (1998).	
	41.	A. Pause et al.; <i>The von Hippel-Lindau tumor-suppressor gene product forms a stable complex with human CUL-2, a member of the Cdc53 family of proteins</i> ; Proc. Natl. Acad. Sci. USA 94 :2156-2161 (1997).	
	42.	F. Ning Li et al.; <i>Grr1 of <i>Saccharomyces cerevisiae</i> is connected to the ubiquitin proteolysis machinery through Skp1: coupling glucose sensing to gene expression and the cell cycle</i> ; The EMBO J. 16 , No. 18:5629-5638 (1997).	
	43.	M. Hochstrasser; <i>Ubiquitin-Dependent Protein Degradation</i> ; Annu. Rev. Genet. 30 :405-439 (1996).	
	44.	P. Kaiser et al.; <i>Cdc34 and the F-box protein Met30 are required for degradation of the Cdk-inhibitory kinase Swe1</i> ; Genes & Dev. 12 :2587-2597 (1998).	
↓	45.	G. Fang et al.; <i>Direct Binding of CDC20 Protein Family Members Activates the Anaphase-Promoting Complex in Mitosis and G₁</i> ; Molecular Cell 2 :163-171.	
DR	46.	H. Yu; <i>Identification of a novel ubiquitin-conjugating enzyme involved in mitotic cyclin degradation</i> ; Current Biology 6 , No 4:455-466.	

EXAMINER
EXAMINER



DATE CONSIDERED 4/25/03

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 U.S. Department of Commerce Patent and Trademark Office				Attorney Docket Number 5470-255			Serial No. 09/541,462
LIST OF DOCUMENTS CITED BY APPLICANT (Use several sheets if necessary)				O P E S C T O P A T E N T & T R A D E M A R K C E N T E R			DEC 08 2000
				Applicants: Xiong et al.			
				Filing Date March 31, 2000			Group 1643-1652
U. S. PATENT DOCUMENTS							
Examiner Initial		Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate
FOREIGN PATENT DOCUMENTS							
		Document Number	Date	Country	Class	Subclass	Translation Yes No
DR	1.	WO 9906554	11/02/99	PCT	C12N	15/12	No
DR	2.	WO 9932514	01/07/99	PCT	C07k	14/00	No
							RECEIVED
DEC 13 2000							
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)							
DR	3.	International Search Report, 11/06/00, for Application No. PCT/US00/08592					TECH CENTER 1600/29
DR	4.	Brown, D. et al. <i>Pip1p, a new subunit of the SCF-Pop ubiquitin ligase complex in S. pombe.</i> Accession No. O13959. 1 June 1998. Database on-line. Available from EMBL.					
DR	5.	Brown, D. et al. <i>S.pombe chromosome I cosmid c23H4.</i> Accession No. Z98977. 8 September 1997. Database on-line. Available from EMBL.					
DR	6.	Fujiwara, T. et al. <i>Human fetal brain cDNA 5'-end GEN-090E07.</i> Accession No. D52876. 24 August 1995. Database on-line. Available from EMBL.					
DR	7.	<i>Hangjun, D. et al. SAG, a novel zinc RING finger protein that protects cells from apoptosis induced by redox agents, Molecular and Cellular Biology.</i> 19:3145-3155 (1999).					
DR	8.	Hillier et al. <i>The WashU-Mrck EST Project.</i> Accession No. W38711. 17 May 1996. Database on-line. Available from EMBL.					
DR	9.	<i>Ohta, T. et al. ROC1, a homolog of APC11, represents a family of cullin partners with an associated ubiquitin ligase activity, Molecular Cell.</i> 3:535-541 (1999).					
DR	10.	<i>Skowyra, D. et al. Reconstitution of G1 Cyclin Ubiquitination with Complexes Containing SCF^{Grl} and Rbx1, Science.</i> 284:662-665 (1999).					
DR	11.	<i>Tanamura, S. et al. MDM2 interacts with MDMX through their RING finger domains, FEBS Letters.</i> 447:5-9 (1999).					

EXAMINER
EXAMINER*BL*

DATE CONSIDERED

4/25/03

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

EXAMINER
*EXAMINER

20 DATE CONSIDERED 4/25/03
Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in
conformance and not considered. Include copy of this form with next communication to applicant